



MA8491-NUMERICAL METHODS
UNIT-I-SOLUTION OF EQUATIONS AND EIGENVALUE PROBLEMS

1. If $g(x)$ is continuous in $[a,b]$, then under what condition the iterative (or) iteration method $x = g(x)$ has a unique solution in $[a,b]$?

- A. $|\phi'(x)| = 1$
- B. $|\phi'(x)| < 1$
- C. $|\phi'(x)| > 1$
- D. $|\phi'(x)| \leq 1$

Answer: (B)

2. What is the other name for iteration method?

- A. Direct method
- B. Successive approximation method**
- C. Power method
- D. Jacobi method

Answer: (B)

3. What is the order of convergence of fixed point iteration method?

- A. 1**
- B. 2
- C. 3
- D. 4

Answer: (A)

4. The formula for iteration method is

- A. $x_{n+1} = \phi(x_n)$**
- B. $x_{n-1} = \phi(x_n)$
- C. $x_n = \phi(x_n)$
- D. $x_n = \phi(x_{n+1})$

Answer: (A)

5. The condition for the convergence of Newton -Raphson method is

- A. $|f(x)f''(x)| = |f'(x)|^2$
 B. $|f(x)f''(x)| = |f(x)|^2$
 C. $|f(x)f''(x)| < |f'(x)|^2$
 D. $|f(x)f''(x)| \leq |f'(x)|^2$

Answer: (C)

6. What is the order of convergence for Newton-Raphson method?

- A. 1
 B. 2
 C. 3
 D. 4

Answer: (B)

7. Newton-Raphson method is also known as

- A. Gauss Elimination method
B. method of Tangents
 C. Gauss –Jacobi method
 D. Power method

Answer: (B)

8. The iterative formula to find \sqrt{N} using Newton-Raphson method is

- A. $x_{n+1} = \frac{x_n^2 + N}{x_n}$
 B. $x_n = \frac{x_n^2 + N}{2x_n}$
 C. $x_{n+1} = \frac{x_n^2 - N}{2x_n}$
 D. $x_{n+1} = \frac{x_n^2 + N}{2x_n}$

Answer: (D)

9. The iterative formula to find cube root of N using Newton-Raphson method is

- A. $x_{n+1} = \frac{x_n^3 + N}{3x_n^2}$
 B. $x_{n+1} = \frac{2x_n^3 + N}{x_n^2}$

$$\text{C. } x_{n+1} = \frac{2x_n^3 + N}{3x_n^2}$$

$$\text{D. } x_{n+1} = \frac{3x_n^3 + N}{2x_n^2}$$

Answer: (C)

10. The iterative formula to find the reciprocal of a positive number using Newton-Raphson method is

$$\text{A. } x_{n+1} = x_n(3 - Nx_n)$$

$$\text{B. } x_{n+1} = x_n(2 - Nx_n)$$

$$\text{C. } x_{n+1} = x_n(2 + Nx_n)$$

$$\text{D. } x_{n+1} = x(2 - Nx_n)$$

Answer: (B)

11. The indirect methods to solve the system of equations is

A. Gauss Elimination

B. Gauss Jordan

C. Crouts

D. Gauss seidel

Answer: (D)

12. From the following which method is called as iterative methods

A. Gauss Elimination method

B. Gauss Jordan method

C. Gauss –Jacobi method

D. Power method

Answer: (C)

13. What type of solutions can be get by using direct methods

A. exact value

B. approximate value

C. moderate value

D. positive value

Answer: (A)

14. What type of solutions can be get by using indirect methods

A. exact value

B. approximate value

C. moderate value

D. positive value

Answer: (B)

15. In Gauss Elimination method the coefficient matrix is reduced to

- A. diagonal matrix
- B. singular matrix
- C. lower triangular matrix
- D. upper triangular matrix**

Answer: (D)

16. In Gauss Jordan method the coefficient matrix is reduced to

- A. diagonal matrix**
- B. singular matrix
- C. lower triangular matrix
- D. upper triangular matrix

Answer: (A)

17. If in each equation of the given system, the absolute value of the largest coefficient is greater than the sum of the absolute values of all the remaining coefficients is called

- A. diagonally dominant**
- B. dominant
- C. absolute value
- D. minimum value

Answer: (A)

18. Solve $x + y = 2$; $2x + 3y = 5$ by Gauss Elimination method.

- A. (1,1)**
- B. (1,2)
- C. (2,1)
- D. (1,-1)

Answer: (A)

19. Solve $3x + 2y = 4$; $2x - 3y = 7$ by Gauss Jordan method.

- A. (1,1)
- B. (1,2)
- C. (2,1)**
- D. (1,-1)

Answer: (C)

20. Solve $x - 2y = 0$; $2x + y = 5$ by Gauss Elimination method.

A. (1,1)

B. (1,2)

C. **(2,1)**

D. (1,-1)

Answer: (C)

21. Solve $2x + y = 3$; $7x - 3y = 4$ by Gauss Jordan method.

A. (1,1)

B. (1,2)

C. (2,1)

D. (1,-1)

Answer: (A)

22. Gauss-Seidel method is twice faster than -----method.

A. Gauss Elimination method

B. Gauss Jordan method

C. Gauss –Jacobi method

D. Power method

Answer: (C)

23. Gauss-Seidel method is better than Gauss –Jacobi method?

A. True

B. False

Answer: (A)

24. What type of eigenvalue can be obtained using power method?

A. simple

B. negative

C. medium

D. dominant

Answer: (D)

25. For what type of matrices, Jacobi's method can be used to find eigen values and eigenvectors?

A. non- symmetric

B. diagonal

C. singular

D. symmetric

Answer: (D)